arXiv:0810.3374v2 [math.NT] 26 Nov 2008

# ON CORRESPONDENCE BETWEEN SOLUTIONS OF A PARAMETRIC FAMILY OF CUBIC THUE EQUATIONS AND ISOMORPHIC SIMPLEST CUBIC FIELDS

AKINARI HOSHI

Abstract. We give a correspondence between non-trivial solutions to a parametric family of cubic Thue equations

$$X^3 - mX^2Y - (m+3)XY^2 - Y^3 = k$$

where $k \mid m^2+3m+9$ and isomorphic simplest cubic fields. By applying R. Okazaki's result for isomorphic simplest cubic fields, we obtain all solutions to the family of cubic Thue equations for $k \mid m^2+3m+9$.

## 1. Introduction

We consider the following parametric family of Thue equations

$$(*) \qquad F_m(X,Y) := X^3 - mX^2Y - (m+3)XY^2 - Y^3 = k$$

for integer $m$ and nonzero integer $k$. We may assume that $-1 \leq m$ and $k > 0$ because if $(x,y)$ is a solution to (*) then $(-x,-y)$ becomes a solution to $F_m(X,Y) = -k$ and $(-y,-x)$ becomes a solution to $F_{-m-3}(X,Y) = k$. For cubic integer $k = a^3$, the equation (*) has three solutions

$$(a,0), \quad (0,-a), \quad (-a,a).$$

We call solutions $(x,y)$ to (*) with $xy(x+y) = 0$ the trivial solutions.

In the case of $k = \pm 1$, Thomas [Tho90] showed that for $n := m+1 \geq 1.365 \times 10^7$ the equation $F_m(X,Y) = \pm 1$ has only the trivial solutions $(\pm 1, 0)$, $(0, \mp 1)$, $(\mp 1, \pm 1)$ and for $0 \leq n = m+1 \leq 10^3$ non-trivial solutions exist only for $m = -1, 0, 2$ (i.e. $n = 0, 1, 3$). Mignotte [Mig93] solved completely the equations $F_m(X,Y) = \pm 1$. He proved that for $-1 \leq m$ non-trivial solutions occur only for $m = -1, 0, 2$.

For $-1 \leq m$ and $k = 1$, all solutions to $F_m(X,Y) = 1$ are given by trivial solutions $(x,y) = (1,0)$, $(0,-1)$, $(-1,1)$ for an arbitrary $m$ and additionally

$$\begin{aligned} &(x,y) = (-1,2),(2,-1),(-1,-1),(4,-9),(-9,5),(5,4) && \text{for} \quad m=-1,\\ &(x,y) = (1,-3),(-3,2),(2,1) && \text{for} \quad m=0,\\ &(x,y) = (-2,9),(9,-7),(-7,-2) && \text{for} \quad m=2 \end{aligned}$$

(cf. also the text book [Gaa02, p.54]). Note that if $(x,y)$ is a solution to (*) then $(y,-x-y)$ and $(-x-y,x)$ are also solutions to (*) because $F_m(X,Y)$ is invariant under the action of the cyclic group $C_3 = \langle \sigma \rangle$ of order three where $\sigma : X \longmapsto Y \longmapsto -X-Y$.

Mignotte-Pethő-Lemmermeyer [MPL96] studied the equation (*) for general $k \in \mathbb{Z}$ and gave a complete solution to Thue inequality $|F_m(X,Y)| \leq 2m+3 = 2n+1$ with a result of

---

2000 *Mathematics Subject Classification.* Primary 11D25, 11D59, 11R16, 12E25.

[LP95]. For $-1 \leq m$ and $1 < k \leq 2m+3$, all solutions to (*) are given by trivial solutions for $k = a^3$ and

$$(x,y) \in \{(-1,2),(2,-1),(-1,-1),(-1,m+2),(m+2,-m-1),(-m-1,-1)\}$$

for $k = 2m+3$, except for $m = 1$ in which case (*) has the extra solutions:

$$(x,y) \in \{(1,-4),(-4,3),(3,1),(3,-11),(-11,8),(8,3)\}$$

for $k = 5$ [1]. Lettl-Pethö-Voutier [LPV99] also investigated Thue inequality $|F_m(X,Y)| \leq k(m)$ where $k : \mathbb{Z} \to \mathbb{N}$ by using the hypergeometric method.

Let $\mathbb{Q}(t)$ be the rational function field over $\mathbb{Q}$ with variable $t$. We take a polynomial

$$f_t^{C_3}(X) = X^3 - tX^2 - (t+3)X - 1 \in \mathbb{Q}(t)[X]$$

which is generic for $C_3$ over $\mathbb{Q}$ in the sense of [JLY02]. For $t = m \in \mathbb{Z}$, the splitting fields $L_m := \mathrm{Spl}_{\mathbb{Q}} f_m^{C_3}(X)$ of $f_m^{C_3}(X)$ over $\mathbb{Q}$ are called simplest cubic fields (cf. [Sha74], [LP95]).

Note that $f_m^{C_3}(X) = F_m(X,1)$ and $F_m(X,1) = F_{-m-3}(-1,-X)$. Thus if $x$ is a root of $f_m^{C_3}(X)$ then $1/x$ becomes a root of $f_{-m-3}^{C_3}(X)$. Hence $L_m = L_{-m-3}$ for $m \in \mathbb{Z}$.

The aim of this note is to give the following theorem which asserts a correspondence between some solutions to (*) and isomorphic simplest cubic fields.

**Theorem 1.** *Let $m$ be an integer. There exists an integer $m' \in \mathbb{Z} \setminus \{m, -m-3\}$ such that $L_m = L_{m'}$ if and only if there exists a solution $(x,y) \in \mathbb{Z}^2$ with $xy(x+y) \neq 0$ to cubic Thue equation*

$$(**) \qquad F_m(x,y) = k \quad \text{for some} \quad k > 0 \quad \text{with} \quad k \mid m^2 + 3m + 9.$$

*Moreover integers $m$, $m'$ and solutions $(x,y)$ to (**) satisfy*

$$m' = m + \frac{(m^2+3m+9)xy(x+y)}{F_m(x,y)} \quad \text{or} \quad -m'-3 = m + \frac{(m^2+3m+9)xy(x+y)}{F_m(x,y)}.$$

Note that (i) the discriminant of $f_m^{C_3}(X)$ with respect to $X$ equals $(m^2+3m+9)^2$, and (ii) if $(x,y)$ is a solution to (**) then $(-x,-y)$ is a solution to $F_m(x,y) = -k < 0$.

In [HM], we showed that for a fixed $m \in \mathbb{Z}$ there exist only finitely many integers $m' \in \mathbb{Z}$ such that $L_m = L_{m'}$. In the range $-1 \leq m < m' \leq 10^5$, we checked with the aid of computer that $L_m = L_{m'}$ if and only if $(m,m') \in \{(-1,5),\ (-1,1259),\ (0,54),\ (5,1259)\} \cup \{(-1,12),\ (0,3),\ (1,66),\ (2,2389),\ (3,54),\ (5,12),\ (12,1259)\}$ (see [HM, Example 5.3]).

Okazaki [Oka02] investigated Thue equations $f(X,Y) = 1$ for irreducible cubic forms $f(X,Y)$ with positive discriminant and established a very strong result on gaps between solutions. Using this gap principle, Okazaki also showed the following theorem:

**Theorem 2** (Okazaki). *For integers $m$, $m' \geq -1$, if $L_m = L_{m'}$ $(m \neq m')$ then $m, m' \in \{-1,0,1,2,3,5,12,54,66,1259,2389\}$. In particular,*

$$L_{-1} = L_5 = L_{12} = L_{1259}, \quad L_0 = L_3 = L_{54}, \quad L_1 = L_{66}, \quad L_2 = L_{2389}.$$

Although a proof of Theorem 2 seems to be unpublished yet (up to now), a brief sketch of the proof is available at `http://www1.doshisha.ac.jp/~rokazaki/papers.html` as a presentation sheet (cf. also [Wak07, §1]).

As a consequence of Theorem 1 and Theorem 2 we obtain the following corollaries:

---

[1] In [MPL96, Theorem 3], there is a misprint: it should be added $(-11,8)$.

**Corollary 3.** *For $m \geq -1$, all solutions $(x, y) \in \mathbb{Z}^2$ with $xy(x+y) \neq 0$ of Thue equations $F_m(X, Y) = k$ with $k > 0$ and $k \mid m^2 + 3m + 9$ are given on Table 1.*

Table 1

| $m$ | $m'$ | $-m'-3$ | $2m+3$ | $k$ | $m^2+3m+9$ | $xy(x+y)$ | $(x,y)$ |
|---|---|---|---|---|---|---|---|
| $-1$ | $-15$ | $12$ | $1$ | $1$ | $7$ | $-2$ | $(-1,2)$, $(2,-1)$, $(-1,-1)$ |
| $-1$ | $1259$ | $-1262$ | $1$ | $1$ | $7$ | $180$ | $(4,-9)$, $(-9,5)$, $(5,4)$ |
| $-1$ | $5$ | $-8$ | $1$ | $7$ | $7$ | $6$ | $(1,-3)$, $(-3,2)$, $(2,1)$ |
| $0$ | $54$ | $-57$ | $3$ | $1$ | $9$ | $6$ | $(1,-3)$, $(-3,2)$, $(2,1)$ |
| $0$ | $-6$ | $3$ | $3$ | $3$ | $9$ | $-2$ | $(-1,2)$, $(2,-1)$, $(-1,-1)$ |
| $1$ | $-69$ | $66$ | $5$ | $13$ | $13$ | $-70$ | $(-2,7)$, $(7,-5)$, $(-5,-2)$ |
| $2$ | $-2392$ | $2389$ | $7$ | $1$ | $19$ | $-126$ | $(-2,9)$, $(9,-7)$, $(-7,-2)$ |
| $3$ | $-3$ | $0$ | $9$ | $9$ | $27$ | $-2$ | $(-1,2)$, $(2,-1)$, $(-1,-1)$ |
| $3$ | $-57$ | $54$ | $9$ | $9$ | $27$ | $-20$ | $(-1,5)$, $(5,-4)$, $(-4,-1)$ |
| $5$ | $1259$ | $-1262$ | $13$ | $49$ | $49$ | $1254$ | $(3,-22)$, $(-22,19)$, $(19,3)$ |
| $5$ | $-15$ | $12$ | $13$ | $49$ | $49$ | $-20$ | $(-1,5)$, $(5,-4)$, $(-4,-1)$ |
| $5$ | $-1$ | $-2$ | $13$ | $49$ | $49$ | $-2$ | $(-1,-2)$, $(-2,3)$, $(3,-1)$ |
| $12$ | $-2$ | $-1$ | $27$ | $27$ | $189$ | $-2$ | $(-1,2)$, $(2,-1)$, $(-1,-1)$ |
| $12$ | $-1262$ | $1259$ | $27$ | $27$ | $189$ | $-182$ | $(-1,14)$, $(14,-13)$, $(-13,-1)$ |
| $12$ | $-8$ | $5$ | $27$ | $189 = 3^3 \cdot 7$ | $189$ | $-20$ | $(-1,5)$, $(5,-4)$, $(-4,-1)$ |
| $54$ | $0$ | $-3$ | $111$ | $343 = 7^3$ | $3087$ | $-6$ | $(-1,-2)$, $(-2,3)$, $(3,-1)$ |
| $54$ | $-6$ | $3$ | $111$ | $1029 = 3 \cdot 7^3$ | $3087$ | $-20$ | $(-1,5)$, $(5,-4)$, $(-4,-1)$ |
| $66$ | $-4$ | $1$ | $135$ | $4563 = 3^3 \cdot 13^2$ | $4563$ | $-70$ | $(-2,7)$, $(7,-5)$, $(-5,-2)$ |
| $1259$ | $-1$ | $-2$ | $2521$ | $226981 = 61^3$ | $1588867$ | $-180$ | $(-4,-5)$, $(-5,9)$, $(9,-4)$ |
| $1259$ | $-15$ | $12$ | $2521$ | $226981 = 61^3$ | $1588867$ | $-182$ | $(-1,14)$, $(14,-13)$, $(-13,-1)$ |
| $1259$ | $5$ | $-8$ | $2521$ | $1588867 = 7 \cdot 61^3$ | $1588867$ | $-1254$ | $(-3,-19)$, $(-19,22)$, $(22,-3)$ |
| $2389$ | $-5$ | $2$ | $4781$ | $300763 = 67^3$ | $5714497$ | $-126$ | $(-2,9)$, $(9,-7)$, $(-7,-2)$ |

**Corollary 4.** *For $m \geq -1$, solutions $(x, y) \in \mathbb{Z}^2$ with $xy(x+y) \neq 0$ to a family of Thue equations $F_m(X, Y) = m^2 + 3m + 9$ exist only for $m = -1, 1, 5, 12, 66, 1259$ as on Table 1.*

We remark that if we assume Okazaki's result (Theorem 2) then by applying Theorem 1 to given $m$ and $m'$, we may get all solutions to (**) without computer.

## 2. PROOF OF THEOREM 1

In [HM], by using formal Tschirnhausen transformation, we gave the following theorem as an analogue to the results of Morton [Mor94] and Chapman [Cha96].

**Theorem 5** (Morton [Mor94], Chapman [Cha96], [HM])**.** *Let $K$ be a field of char $K \neq 2$. For $m, m' \in K$, the splitting fields of $f_m^{C_3}(X)$ and of $f_{m'}^{C_3}(X)$ over $K$ coincide if and only if there exists $z \in K$ such that either*

$$m' = \frac{m(z^3 - 3z - 1) - 9z(z+1)}{mz(z+1) + z^3 + 3z^2 - 1} \quad \text{or} \quad m' = -\frac{m(z^3 + 3z^2 - 1) + 3(z^3 - 3z - 1)}{mz(z+1) + z^3 + 3z^2 - 1}.$$

We use Theorem 5 in the case where $K = \mathbb{Q}$.

Assume that there exists an integer $m' \in \mathbb{Z} \backslash \{m, -m-3\}$ such that $L_m = L_{m'}$. By Theorem 5, there exists $z \in \mathbb{Q}$ such that either

$$(1) \qquad m' = \frac{m(z^3 - 3z - 1) - 9z(z+1)}{mz(z+1) + z^3 + 3z^2 - 1} \quad \text{or} \quad m' = -\frac{m(z^3 + 3z^2 - 1) + 3(z^3 - 3z - 1)}{mz(z+1) + z^3 + 3z^2 - 1}.$$

Note that $mz(z+1)+z^3+3z^2-1=z^3+(m+3)z^2+mz-1=-F_m(1,z)$. If $z=0$ then $m'=m$ or $m'=-m-3$. Hence it follows $z\neq 0$. We put

$$u:=\frac{1}{z}$$

then the condition (1) can be written as

$$m'=m+\frac{(m^2+3m+9)u(u+1)}{F_m(u,1)}\quad \text{or}\quad m'=-m-3-\frac{(m^2+3m+9)u(u+1)}{F_m(u,1)}.$$

Write $u=x/y$ as a quotient of relatively prime integers $x,y$. Then there exist relatively prime $x,y\in\mathbb{Z}$ such that

$$m'=m+\frac{(m^2+3m+9)xy(x+y)}{F_m(x,y)}\quad \text{or}\quad -m'-3=m+\frac{(m^2+3m+9)xy(x+y)}{F_m(x,y)}.$$

In particular, we have

$$\frac{(m^2+3m+9)xy(x+y)}{F_m(x,y)}\in\mathbb{Z}. \tag{2}$$

We will show that $k:=F_m(x,y)$ divides $(m^2+3m+9)$.

Put $h(u):=(m^2+3m+9)u(u+1)$ and $f(u):=F_m(u,1)$. We take the resultant $R:=\mathrm{Res}_u(h(u),f(u))$ of $h(u)$ and $f(u)$ with respect to $u$. From the definition we have

$$R=\begin{vmatrix} m^2+3m+9 & m^2+3m+9 & 0 & 0 & 0\\ 0 & m^2+3m+9 & m^2+3m+9 & 0 & 0\\ 0 & 0 & m^2+3m+9 & m^2+3m+9 & 0\\ 1 & m+3 & m & -1 & 0\\ 0 & 1 & m+3 & m & -1 \end{vmatrix}=-(m^2+3m+9)^3. \tag{3}$$

We see $R$ is also given of the form

$$\begin{aligned} R&=\begin{vmatrix} m^2+3m+9 & m^2+3m+9 & 0 & 0 & h(u)u^2\\ 0 & m^2+3m+9 & m^2+3m+9 & 0 & h(u)u\\ 0 & 0 & m^2+3m+9 & m^2+3m+9 & h(u)\\ 1 & m+3 & m & -1 & f(u)u\\ 0 & 1 & m+3 & m & f(u) \end{vmatrix}\\ &=-(m^2+3m+9)^2\Big(h(u)p(u)+f(u)q(u)\Big) \end{aligned} \tag{4}$$

where

$$p(u)=2u^2-2mu-u-m-5,\quad q(u)=-(m^2+3m+9)(2u+1).$$

By (3) and (4), we have

$$h(u)p(u)+f(u)q(u)=(m^2+3m+9).$$

Put

$$\begin{aligned} H(x,y)&:=(m^2+3m+9)xy(x+y),\\ P(x,y)&:=2x^2-2mxy-xy-my^2-5y^2,\\ Q(x,y)&:=-(m^2+3m+9)y(2x+y). \end{aligned}$$

Then it follows from $u = x/y$ that

$$H(x,y)P(x,y) + F_m(x,y)Q(x,y) = (m^2+3m+9)y^5.$$

Because the cubic forms $F_m(X,Y)$ and $H(X,Y)$ are invariants under the action of $\sigma^2$ : $Y \longmapsto X \longmapsto -X-Y$, we also get

$$H(x,y)P(-x-y,x) + F_m(x,y)Q(-x-y,x) = (m^2+3m+9)x^5.$$

Hence by (2) we have

$$\begin{aligned}\frac{H(x,y)P(x,y)}{F_m(x,y)} + Q(x,y) &= \frac{(m^2+3m+9)y^5}{F_m(x,y)} \in \mathbb{Z},\\ \frac{H(x,y)P(-x-y,x)}{F_m(x,y)} + Q(-x-y,x) &= \frac{(m^2+3m+9)x^5}{F_m(x,y)} \in \mathbb{Z}.\end{aligned}$$

Since $x$ and $y$ are relatively prime, we conclude that $k = F_m(x,y)$ divides $(m^2+3m+9)$.

Conversely if there exists relatively prime $x, y \in \mathbb{Z}$ such that $k = F_m(x,y)$ divides $(m^2+3m+9)$ then we can take $m' \in \mathbb{Z}$ as

$$m' = m + \frac{(m^2+3m+9)xy(x+y)}{F_m(x,y)} \quad \text{or} \quad -m'-3 = m + \frac{(m^2+3m+9)xy(x+y)}{F_m(x,y)}.$$

Note that $m = m'$ or $m = -m'-3$ if and only if $xy(x+y) = 0$. ☐

**Acknowledgment.** This work is partially supported by Rikkyo University Special Fund for Research.

## References

[Cha96] R. J. Chapman, *Automorphism polynomials in cyclic cubic extensions*, J. Number Theory **61** (1996), 283–291.

[Gaa02] I. Gaál, *Diophantine equations and power integral bases. New computational methods*, Birkhäuser Boston, Inc., Boston, MA, 2002.

[HS00] M. Hindry, J. H. Silverman, *Diophantine geometry. An introduction*, Graduate Texts in Mathematics, 201, Springer-Verlag, New York, 2000.

[HM07] A. Hoshi, K. Miyake, *Tschirnhausen transformation of a cubic generic polynomial and a 2-dimensional involutive Cremona transformation*, Proc. Japan Acad. Ser. A **83** (2007), 21–26.

[HM] A. Hoshi, K. Miyake, *A geometric framework for the subfield problem of generic polynomials via Tschirnhausen transformation*, preprint. arXiv:0710.0287v2 [math.NT]

[JLY02] C. Jensen, A. Ledet, N. Yui, *Generic polynomials, constructive aspects of the inverse Galois problem*, Mathematical Sciences Research Institute Publications, Cambridge, 2002.

[Lan78] S. Lang, *Elliptic curves: Diophantine analysis*, Grundlehren der Mathematischen Wissenschaften 231, Springer, 1978.

[Lan83] S. Lang, *Fundamentals of Diophantine geometry*, Springer-Verlag, New York, 1983.

[LP95] F. Lemmermeyer, A. Pethö, *Simplest cubic fields*, Manuscripta Math. **88** (1995), 53–58.

[LPV99] G. Lettl, A. Pethö, P. Voutier, *Simple families of Thue inequalities*, Trans. Amer. Math. Soc. **351** (1999), 1871–1894.

[Mig93] M. Mignotte, *Verification of a conjecture of E. Thomas*, J. Number Theory **44** (1993), 172–177.

[MPL96] M. Mignotte, A. Pethö, F. Lemmermeyer, *On the family of Thue equations* $x^3-(n-1)x^2y-(n+2)xy^2-y^3=k$, Acta Arith. **76** (1996), 245–269.

[Mor94] P. Morton, *Characterizing cyclic cubic extensions by automorphism polynomials*, J. Number Theory **49** (1994), 183–208.

[Oka02] R. Okazaki, *Geometry of a cubic Thue equation*, Publ. Math. Debrecen **61** (2002), 267–314.

[Sha74] D. Shanks, *The simplest cubic fields*, Math. Comp. **28** (1974), 1137–1152.

[Sie29] C. L. Siegel, *Über einege Anwendungen Diophantischer Approximationen*, Abh. Preuss. Akad. Wiss. Phys. Math. Kl. (1929), 41–69.

[Tho90] E. Thomas, *Complete solutions to a family of cubic Diophantine equations*, J. Number Theory **34** (1990), 235–250.

[Wak07] I. Wakabayashi, *Number of solutions for cubic Thue equations with automorphisms*, Ramanujan J. (2007), 131–154.

Akinari Hoshi
Department of Mathematics
Rikkyo University
3–34–1 Nishi-Ikebukuro Toshima-ku
Tokyo, 171–8501, Japan
E-mail: `hoshi@rikkyo.ac.jp`
Web: `http://www2.rikkyo.ac.jp/web/hoshi/`